

Two-Dimensional Electrophoresis of Human Plasma Apolipoproteins

Dennis L. Sprecher, Lila Taam, and H. Bryan Brewer, Jr.

The two-dimensional electrophoretic method with silver staining we describe better resolves plasma apolipoproteins (apo) than any procedure previously described. It can be used to screen for abnormalities in apoA-I, apoA-II, apoA-IV, apoC-II, apoC-III, apoD, apoE, and apoH. In addition, this is the first presentation of apoD and apoH on two-dimensional gels. This electrophoretic method will permit the quantification of plasma apolipoproteins by computer imaging. Such detailed analysis of the plasma apolipoproteins should provide a better understanding of apolipoprotein function and the role these apolipoproteins play in lipoprotein metabolism and the pathophysiology of dyslipoproteinemias and coronary artery disease.

Additional Keyphrases: heritable disorders · screening · atherosclerosis · heart disease · dyslipoproteinemia · apolipoproteins D and H

Apolipoproteins are the protein moieties of lipoprotein particles. Over the last several years apolipoproteins have been extensively investigated and have been shown to modulate the activity of enzymes that are important in lipid metabolism (e.g., lipoprotein lipase, lecithin-cholesterol acyltransferase) and to play a pivotal role, on lipoprotein particles, as the ligand for receptor-lipoprotein interactions (1). In addition, apolipoprotein concentrations both in plasma and in separated lipoprotein fractions have been found to correlate highly with the development of coronary artery disease in certain populations (2, 3).

Familial absences as well as structural defects in plasma apolipoproteins leading to clinical dyslipoproteinemias have now been recognized. Molecular defects in plasma apolipoproteins, therefore, are analogous to the hemoglobinopathies that have previously been shown to be ascribable to amino acid substitutions, insertions in the hemoglobin genes, and defects in post-translational modifications (4).

Structural defects in the plasma apolipoproteins may affect either the charge or the molecular mass of the protein. Two-dimensional electrophoresis, which characterizes proteins by both of these characteristics, may be used to identify up to 30% of the known amino acid substitutions (5).

The purpose of this study is to present a relatively simple two-dimensional electrophoretic method for evaluating plasma apolipoproteins A-I (apoA-I), A-II, A-IV, C-II, C-III, D, E, and H.¹ In addition, apoC-I may be evaluated by using an extended pH range and apoB-48 as well as apoB-100 by use of 3% polyacrylamide SDS gel electrophoresis. We find this

technique effective in screening for molecular defects in the plasma apolipoproteins in patients with dyslipoproteinemias and atherosclerosis.

Over the years, various nomenclatures have been used to identify the plasma lipoproteins that are resolved on one-dimensional and two-dimensional gel electrophoresis (6-8). Inconsistency and lack of a standardized nomenclature for the plasma apolipoproteins have resulted in confusion in the literature. Therefore, we have adopted a nomenclature for the apolipoproteins that is consistent with the nomenclature currently standardized for identification of proteins separated by two-dimensional electrophoresis (9-11).

Materials and Methods

Samples. Blood was collected in tubes containing Na₂EDTA (final concentration, about 1 g/L) as anticoagulant. The erythrocytes were removed and the plasma was stored at 4 °C for as long as two weeks before analysis.

For two-dimensional electrophoresis we used plasma or isolated lymph chylomicrons, VLDL, and HDL. Human thoracic duct lymph was collected and chylomicrons were separated by centrifugation in an SW 27 Beckman swinging-bucket rotor for 115 min at 25 000 rpm and 4 °C. VLDL was isolated by ultracentrifugation (12) of 5 mL of plasma in a 40.3 Beckman rotor (Beckman Instrument Co., Fullerton, CA) at 39 000 rpm ($\approx 110\,000 \times g$) at 4 °C for 18 h. HDL was isolated by ultracentrifugation between relative densities of 1.063 to 1.21 (4 °C, 36 h, 39 000 rpm).

Lipoprotein fractions were recovered by tube slicing, desalted by chromatography on a 2 × 5.5 cm column of Sephadex G-25M (PD-10; Pharmacia Fine Chemicals, Uppsala, Sweden), equilibrated with an 0.8 g/L solution of ammonium bicarbonate. Lipoproteins were lyophilized and delipidated with chloroform/methanol (2/1 by vol) (13). The protein pellet was dried under a stream of nitrogen, dissolved by stirring in freshly prepared 8 mol/L urea containing 1 mg of DTT per liter, and then incubated for 30 min at 37 °C.

Plasma samples were prepared as reported by O'Farrell (14), with minor modifications. Two microliters of plasma were placed in a "detergent solution," containing, per liter, 20 g of SDS, 50 mL of β -mercaptoethanol, 200 mL of glycerol, 20 mL of Nonidet-P40 detergent, 60 g of CHAPS, and 50 mL of ampholytes (pH 5-7 and 4-6, 1/2 by vol; Serva, Heidelberg, F.R.G.). Stored at -20 °C, this solution was stable for as long as three months. For analysis, we added 10 μ L of detergent solution to 2 μ L of plasma, then 8 mg of solid urea, and followed this with intermittent vortex-mixing for the next 20 min, at room temperature.

For treatment with neuraminidase (EC 3.2.1.18) (15) we mixed 100 μ L of plasma with 100 μ L of a buffer containing 1 U of neuraminidase (from *Clostridia perfringens*; Sigma Chemical Co., St. Louis, MO), and, per liter, 0.1 mmol of potassium monophosphate (pH 4.5) and 1 mmol of calcium chloride. The pH was adjusted to 5.0 with HCl and the mixture was incubated at 37 °C for 4 h. We then added 4 μ L of this solution to 10 μ L of the detergent solution, followed by 9 mg of solid urea.

Molecular Disease Branch, National Heart, Lung, and Blood Institute, Building 10, Room 7N-117, National Institutes of Health, Bethesda, MD 20205.

¹ Nonstandard abbreviations: SDS, sodium dodecyl sulfate; apo, apolipoprotein; VLDL, very-low-density lipoproteins; LDL, low-density lipoproteins; HDL, high-density lipoproteins; DTT, 1,4-dithiothreitol; CHAPS, 3-[(3-cholamidopropyl)dimethyl ammonio]-1-propane sulfonic acid; and LCAT, lecithin-cholesterol acyltransferase (EC 2.3.1.43).

Received July 9, 1984; accepted September 7, 1984.

Two-dimensional gel electrophoresis was performed similarly to the methods described previously (9, 16) with modifications. Polyacrylamide gels (3 g of acrylamide and 88 mg of bisacrylamide dissolved in 38.5 mL of freshly prepared 8.6 mol/L urea) for isoelectrofocusing were prepared in 1.75 × 95 mm glass tubes with 1.5 mL of the above-mentioned ampholytes. Ammonium persulfate (75 μL of an 180 mL/L stock solution) and 40 μL of *N,N,N',N'*-tetramethylethylenediamine (Bio-Rad Lab., Richmond, CA) were used as polymerization catalysts. Gels were polymerized for 30 min, and either the plasma mixture described above or delipidated lymph chylomicrons (35 μg), VLDL (30 μg), or HDL (50 μg) redissolved in urea and DTT was placed on the gel and carefully overlaid with NaOH, 20 mmol/L. Solutions in the upper tray and the lower tray were 20 mmol/L NaOH and 10 mmol/L H₃PO₄, respectively. For plasma, the gels were electrofocused (at 4 °C) at 250 V for the first 2 h, 500 V for the next 15 h, and 1000 V for the last 2 h, a total of 10 000 V · h. Isolated delipidated lipoprotein fractions were electrofocused at 250 V for 16 h at 4 °C.

After isoelectrofocusing, gels were either rapidly frozen on solid CO₂ and stored (for as long as a month) or placed for 10 min at room temperature in an equilibration solution containing (per liter) 0.1 mol of Tris (pH 6.8), 10 g of SDS, and 25 mL of β-mercaptoethanol. The gels were then loaded on the top of a 11 cm × 12 cm × 0.75-mm-thick polyacrylamide slab gel (per liter, 150 g of acrylamide, 1 g of bisacrylamide, and 1 g of SDS). A stacking gel (4.0% T, 2.7% C) was used for the analysis of plasma samples. The gels were electrophoresed until the dye front was 5 mm from the bottom of the gel (approximately 3.5 h, 10 mA per gel for stacking, 20 mA per gel for running gel; 4 °C) and then stained in Coomassie Blue or silver stain. We stained with Coomassie Brilliant Blue R250 (1 g/L) after fixing the gels in acetic acid/methanol/water (5/50/45 by vol). We destained them in 50 mL/L methanol and 75 mL/L acetic acid until a clear background was obtained. The silver stain used for the plasma gels was similar to that reported by Merrill et al. (17) and Morrisey (18).

Immunoblotting was performed as previously described (19). Plasma gels were blotted at 80 V for 1 h onto nitrocellulose sheets. Isolation of antibodies to apoA-II and apoD has been previously reported (20, 21). Second antibody was used as described in manufacturer's (Bio-Rad Lab.) instructions.

Results

Table 1 lists the major human plasma apolipoproteins and some of their physico-chemical properties. Figure 1 shows a gel on which the human plasma apolipoproteins are re-

solved by use of a narrow pH range (4.0–7.0). Figure 2 illustrates an electrophoretogram for a sample preincubated with neuraminidase (EC 3.2.1.18). ApoB does not appear on these gels, owing to its high molecular mass, nor does apoC-I, whose pI is outside the 4–7 pH range.

Figures 3, 4, and 5 illustrate two-dimensional electrophoretograms of the apolipoproteins present in isolated lymph




Fig. 1. Two-dimensional electrophoretogram of human plasma, pH 4–7, with specific apolipoproteins circled




Fig. 2. Two-dimensional electrophoretogram of human plasma treated with neuraminidase, allowing easier identification of apoA-IV, apoE, and the desialylated form of apoD. ApoH migrates off the basic end of the electrophoretogram




Fig. 3. Two-dimensional electrophoretogram of human thoracic duct lymph chylomicrons. ApoH, apoA-IV, and the C apolipoproteins are increased in this density fraction

Table 1. Properties of the Major Apolipoproteins in Human Plasma

| Apolipo-protein | M _r | No. of amino acids | Chromo-some | Concn in plasma, mg/L | pI of major isoform* |
|-----------------|----------------|--------------------|-------------|-----------------------|----------------------|
| A-I | 27 000 | 243 | 11 | 1200 | 5.6 |
| A-II | 17 000 | 154 ^b | 1 | 300 | 4.9 |
| A-IV | 46 000 | 393 | 11 | 40 | 5.5 |
| B-100 | 500 000 | — | — | 900 | — |
| B-48 | 250 000 | — | — | <50 | — |
| C-I | 6500 | 57 | — | 50 | 8.0 |
| C-II | 8800 | 79 | 19 | 50 | 4.7 |
| C-III | 8750 | 79 | 11 | 120 | 4.9 |
| D | 32 000 | — | — | 120 | 5.3 |
| E | 35 000 | 299 | 19 | 50 | 5.9 ^c |
| H | 48 000 | — | — | 70 | 9.5 |

* Designated with zero subscript in text.

^b Dimer.

^c ApoE₀; also has been reported as ApoE-3 (see text).




Fig. 4. Two-dimensional electrophoretogram of VLDL of human plasma. ApoE, apoC-II, and apoC-III are particularly enriched in this density fraction.




Fig. 5. Two-dimensional electrophoretogram of HDL of human plasma. ApoA-I and apoA-II are noted to be increased in this density fraction.

chylomicrons, VLDL, and HDL, respectively. Lymph chylomicrons have a greater concentration of apoH, apoA-IV, and the C apolipoproteins. VLDL has a higher concentration of apoE, apoC-II, and apoC-III; HDL, a greater concentration of apoA-I and apoA-II. Note that albumin, α_1 -antitrypsin, and prealbumin are present in both VLDL and HDL fractions. By comparing Figure 1 with Figures 3, 4, and 5, one can evaluate the total apolipoproteins present in human plasma with those in the individual density fractions.

ApoA-I

ApoA-I, a major HDL apolipoprotein, is biosynthesized as a 267-residue precursor protein, preproapoA-I (22–25), which undergoes co-translational cleavage. The resulting proapoA-I contains a hexapeptide attached to the amino-terminus of mature apoA-I; it is secreted into plasma and lymph (26) and undergoes extracellular post-translational cleavage to the mature 243-residue apoA-I (27). In vitro, apoA-I activates LCAT, the enzyme that catalyzes the esterification of cholesterol in plasma (28). ApoA-I has been shown to be located in the p11–q13 region of human chromosome 11 (29). Decreased concentrations of apoA-I in

plasma have been correlated with increased risk of premature coronary artery disease (30).

Mature apoA-I is polymorphic in plasma (Figure 1). The major isoform in plasma, designated apoA-I₀, has been previously designated in the literature as apoA-I₃ (26), apoA-I₄ (7), or apoA-I₁ (31). ApoA-I₊₂ in plasma and lymph is proapoA-I (Figure 6) (26). The minor isoforms at apoA-I₊₁ and apoA-I₋₁ are primarily de-amidated proapoA-I and mature apoA-I, respectively.

Several apoA-I variants have now been recognized. The most extensively studied is apoA-I_{Tangier} (32, 33), characterized by a marked increase in apoA-I₊₂ or proapoA-I_{Tangier} (Figure 6). Several additional apoA-I variants identified by two-dimensional electrophoresis are summarized in Table 2 (31, 34–36). ApoA-I variants may be associated with subnormal concentrations of HDL and with premature cardiovascular disease.

ApoA-II

ApoA-II, the second major protein of HDL, is biosynthesized as a 100-residue precursor protein, preproapoA-II (37). PreproapoA-II is co-translationally cleaved to proapoA-II, which contains a pentapeptide attached to mature apoA-II (38). Mature apoA-II exists as a 77-residue monomer (apoA-II_{monomer}) and a 154-residue dimer (apoA-II_{dimer}) in which two monomers are linked by a disulfide bridge at position 6 in the sequence (20). ApoA-II has been localized to the p21–qter region of human chromosome 1 (39). No definitive function of apoA-II has been established. ApoA-II activates hepatic lipase in vitro (40), but the physiological importance of this observation remains to be established. ApoA-II concentrations in plasma reportedly are inversely associated with the development of premature cardiovascular disease (41).

Mature apoA-II is also polymorphic in plasma (Figure 1) (42), and the major isoform is designated apoA-II₀. The modifications of apoA-II responsible for the isoforms seen immediately to the acidic and basic sides of apoA-II₀ (Figure 7) are unknown. Analysis of the apoA-II isoforms separated on two-dimensional gel electrophoresis at pH 3.0–10.0 by the immunoblot technique revealed an additional isoform of apoA-II, designated apoA-II₊₃ (Figure 7), now known to be the proform, proapoA-II (43).

ApoA-IV

ApoA-IV, a 393-residue protein, M_r 46 000, is present on triglyceride-rich lipoproteins and may be involved in their metabolism (44, 45). ApoA-IV has been provisionally assigned to chromosome 11 (46). The major apoA-IV isoform,




Fig. 6. Two-dimensional analysis of purified normal apoA-I, which usually has at least five isoforms, and apoA-I_{Tangier}. The mature form is specified as apoA-I₀, and the proisoprotein as apoA-I₊₂. Note that the apoA-I concentration in Tangier disease is decreased, and more proisoprotein is present than mature isoprotein.

Table 2. ApoA-I Variants*

| | (Acidic) | | Apo A-I (pI 5.60) | | | | (Basic) |
|------------------------------------------------------------|------------------------------|----|----------------------|----|----|----|--------------------------|
| | -2 | -1 | 0 | +1 | +2 | +3 | |
| Apo A-I | | • | ● | | • | | |
| Apo A-I _{Milano} (Arg ₁₇₃ → Cys) | • | ● | ● | • | • | | |
| Apo A-I _{Munster-2} (Lys ₁₀₇ deletion) | } | ● | ● | • | • | | |
| Apo A-I _{Marburg} (Lys ₁₀₇ deletion) | | | | | | | |
| Apo A-I _{Giessen} (Pro ₁₄₃ → Arg) | | • | ● | ● | • | • | |
| Apo A-I _{Munster-3} { | } (Asp ₁₀₃ → Asn) | • | ● | ● | • | • | |
| | | | | | | | (Pro ₄ → Arg) |
| | | | | | | | |
| Apo A-I _{Tangier} | | | • | | • | | |
| Apo A-I + Apo C-III Deficiency | | | | | | | |

* ApoA-I variants, indicated above with amino acid substitutions, are all represented symbolically in the table as the reported heterozygote, i.e., products of one normal and one abnormal allele. Isoforms represented by dots to the basic side of the mature form(s), are the proforms two positive charge units away from the corresponding mature form(s).




Fig. 7. Immunoblot of apoA-II, separated on two-dimensional electrophoresis (pH 3-10), revealing multiple isoforms of apoA-II in human plasma

apoA-IV₀, has a pI of 5.50. Two minor isoforms, apoA-IV₋ and apoA-IV₊ (number of charge units away from major form unknown), are present in plasma. In approximately 10% of the German population, the major isoform of apoA-IV migrates as a doublet, with an additional isoform presumably one charge unit to the basic side (Figure 8), and has been designated apoA-IV_{Marburg} (31). Several structural apoA-IV variants have been reported (8), but the significance of these isoforms on apoA-IV function is as yet unknown (Table 3).

ApoB

Human apoB can be separated into two forms, apoB-48 and apoB-100, which are synthesized by the intestine and liver, respectively (47-49). ApoB-100 interacts with the LDL or apoB,E receptor (50). ApoB-48, a major protein of chylomicrons, may function as a structural protein for lipoprotein particles secreted by the intestine. Deficiency of apoB-100 and apoB-48 results in abetalipoproteinemia and homozygous hypobetalipoproteinemia, a condition in which the only lipoprotein in plasma is HDL (51). ApoB-100 and apoB-48 are not visible on two-dimensional electrophoretograms of plasma, owing to their large size. Rather, they are




Fig. 8. Two-dimensional gel electrophoresis of purified apoA-IV. Ninety percent of the general German population have the normal single isoform, 10% have the doublet or apoA-IV_{Marburg} variant

Table 3. ApoA-IV Variants

| | ApoA-IV (pI 5.50) (pI 5.53) | | | |
|------------------------------------------------|--------------------------------|---|----|----|
| | -1 | 0 | +1 | +2 |
| ApoA-IV (90% of population) | | ● | | |
| ApoA-IV _{Marburg} (10% of population) | | ● | ● | |
| ApoA-IV variant | | ● | ● | |
| ApoA-IV variant | | ● | | ● |
| ApoA-IV variant | ● | ● | | |

We assume that these variants are separated by single charge units. Acidic and basic directions as in Table 2.

most readily separated and identified on SDS-agarose gel electrophoresis (Figure 9) (52).

ApoC-I

ApoC-I is synthesized as a 83-residue precursor protein, preapoC-I (53). Mature apoC-I, which contains 57 amino




Fig. 9. A, normal VLDL with apoB-100; B, VLDL isolated from a patient with apoE deficiency, which is enriched in apoB-48

ApoB is too large to be separated by two-dimensional gel electrophoresis and is more readily analyzed by 3% agarose gel/SDS electrophoresis (51) as shown here

acids (54, 55), reportedly inhibits hepatic lipase (36) and activates LCAT (56). It is not seen on the plasma electrophoretograms illustrated in Figure 1 because its pI is 8.0. Figure 10, which shows a gel in which a pH range of 6 to 9 was used, illustrates the relative positions of apoC-I and apoA-I. ApoC-I appears to have two isoforms.

ApoC-II

ApoC-II is a 79-residue apolipoprotein that functions as a co-factor for lipoprotein lipase, the major enzyme catalyzing the hydrolysis of plasma triglycerides (57, 58). ApoC-II is synthesized as a 101-residue precursor, preapoC-II (59), which undergoes co-translational cleavage to mature apoC-II (60, 61). ApoC-II has been localized to chromosome 19 (62, 63).

The major apoC-II isoform, apoC-II₀, has a pI of 4.7. Although a minor isoform (demonstrated here as apoC-II₋₁) has been reported (64), it has not been consistently identified on the plasma electrophoretograms (Figures 1, 11). Immunoblot techniques have further identified two other isoforms for apoC-II, illustrated in Figures 3 and 4.

Patients with a deficiency of apoC-II are characterized clinically by supranormal triglyceride concentrations in their plasma and a type I hyperlipoproteinemia (65, 67). Figure 11 depicts electrophoretograms of plasma from a patient with apoC-II deficiency, which demonstrates a re-




Fig. 10. Two-dimensional electrophoretogram (pH 6-9) of purified apoC-I (pI = 8.0), and its relationship to apoA-I




Fig. 11. Two-dimensional electrophoretograms (left) of apoC-II and apoC-III with and without neuraminidase treatment; right, results of similar analysis and treatment plasma from a patient with apoC-II deficiency

Note that the -1 and -2 isoforms of apoC-III disappear with desialylation. In addition, the apoC-II deficient patient has a low concentration of apoC-II, which migrates to an abnormal electrophoretic position. The dashed circle illustrates the relative electrophoretic position of normal apoC-II

duced level of apoC-II. ApoC-II is identified, however, in an abnormal electrophoretic position.

ApoC-III

ApoC-III is synthesized as a 99-residue protein, preapoC-III (68). This is co-translationally cleaved to yield mature apoC-III, which is present in plasma as three isoforms containing no, one, or two neuraminic acid residues. The major desialylated isoform, apoC-III₀, is a minor isoform in plasma. The two predominant isoforms, apoC-III₋₁ and apoC-III₋₂, contain one and two sialic acid residues, respectively (69) (Figure 11). The pI of the apoC-III₀, apoC-III₋₁, and apoC-III₋₂ isoforms are 4.9, 4.65, and 4.5, respectively. Figure 3 also illustrates additional minor apoC-III isoforms with lower isoelectric points. ApoC-III prevents lipoprotein uptake by the perfused rat liver, which suggests that it may modulate the metabolism of triglyceride-rich lipoproteins (70, 71).

Patients with a deficiency of apoA-I+apoC-III and severe atherosclerosis have been reported (36, 72). Patients with apoA-I+apoC-III deficiency have relatively normal concentrations of LDL in their plasma, but HDL are virtually absent. One kindred of patients with apoA-I+apoC-III deficiency is characterized by an insertion in the apoA-I gene (73). Restriction enzyme analysis of the genomic DNA of a second kindred with apoA-I+apoC-III deficiency has shown no major deletions or insertions in either the apoA-I or apoC-III gene (72).

ApoD

ApoD is a glycoprotein containing 18% carbohydrate by weight. Its M_r is approximately 32 000 (21, 74). ApoD is mostly present on HDL. It may function with LCAT and apoA-I in the esterification of cholesterol and the transfer of cholesteryl esters from HDL to VLDL (75).

ApoD is polymorphic in plasma, the six isoforms being primarily ascribable to carbohydrate heterogeneity (Figure 12). The major isoform of desialylated apoD, apoD₀, has a pI of 5.3 (Figure 2). The electrophoretic relationship of apoA-I, apoA-II, and apoD, as evaluated by the immunoblot technique, is illustrated in Figure 12.




Fig. 12. Immunoblots of two-dimensional electrophoretograms of normal human plasma, indicating the relative positions of apoD, apoA-I, and apoA-II

The pI of the mature, desialylated form, apoD₀, depicted (panel B) is 5.3

ApoE

ApoE, a 299-residue glycoprotein, is synthesized as an amino acid precursor protein, preapoE (76, 77). PreapoE undergoes co-translational cleavage of the 18-residue prepeptide to mature apoE, which is secreted into plasma as a sialylated apolipoprotein. Newly secreted sialylated apoE loses the sialic acid residues, and the predominant apoE isoforms in plasma contain no sialic acid (Figures 1 and 2). ApoE is located at a single genetic locus, and there are three common alleles, designated ϵ_2 , ϵ_3 , ϵ_4 (Figure 13). As a result, six common apoE phenotypes are present in the population: E2/2, E3/3, E4/4, E2/3, E2/4, and E3/4 (6). The common apoE isoforms differ at two positions in the amino acid sequence (i.e., positions 112 and 158), apoE-2 containing Cys 112 and Cys 158, apoE-3 Cys 112 and Arg 158, and apoE-4 Arg 112 and Arg 158. The most common allele, ϵ_3 , is considered to be a normal allele. The locus for apoE may be located on human chromosome 19 (78).

ApoE is also considered important in the metabolism of triglyceride-rich lipoproteins and to bind in vitro to a specific membrane receptor in the liver (79). The E2/2 phenotype is associated with type III hyperlipoproteinemia (80-82). Previous studies have established that the apoE-2 isoprotein isolated from type III patients has delayed catabolism (81) and defective binding to the cellular apoE receptor (82). The several additional structural variants of apoE reported are summarized in Table 4 (16, 83-90). A recently identified kindred with phenotype E1/3 is also presented (Figure 13).

The nomenclature of apoE has been confusing and has recently been reviewed (6). In the present report, we locate apoE-3, the predominant normal plasma isoform, at the designated apoE₀ position, for consistency with the nomen-




Fig. 13. Two-dimensional electrophoretograms of apoE isoproteins relative to the electrophoretic position of apoA-I: upper and lower panels, common apoE isoproteins; middle panel, rare apoE₋₂ isoprotein, present in greater concentration than the common apoE₀ isoprotein

clature for proteins separated by two-dimensional electrophoresis. The other two major isoforms, apoE-2 and apoE-4 are located at positions designated apoE₋₁ and apoE₊₁, respectively. However, the previous apoE-2, apoE-3, and apoE-4 designations are also included in parentheses in the two-dimensional electrophoretograms in Figure 12. The importance of the standardized nomenclature for apolipoproteins separated by electrophoresis is particularly relevant to the categorization of apoE isoforms, which have charge differences based on amino acid substitutions as well as post-translational modifications. Many variants, therefore, may have a similar electrophoretic position on two-dimensional gel electrophoresis, but may not necessarily be the identical protein; e.g., an apoE₋₁ isoform may be apoE-2 or a de-amidated form of apoE-3. Therefore, two different structural apoE isoforms may be located at the same electrophoretic position, and amino acid sequence analysis may be required to differentiate the two different isoproteins. The potential for different E apolipoproteins to be located at the same electrophoretic position illustrates the need for the

Table 4. ApoE Variants

| | ApoE | | | | | (Basic) |
|----------------------------------------------------------------------------------|----------|----------|----------|----------|----------|---------|
| | (Acidic) | -2 | -1 | 0 | +1 | |
| | (ApoE-1) | (ApoE-2) | (ApoE-3) | (ApoE-4) | (ApoE-5) | |
| ApoE ₀ (ApoE-3-normal isoprotein) | | | • | | | |
| ApoE ₊₁ [ApoE-4 (Cys ₁₁₂ → Arg)] | | | | • | | |
| ApoE ₋₁ [ApoE-2 (Arg ₁₅₈ → Cys)] | | • | | | | |
| ApoE ₋₁ [ApoE-2 (Arg ₁₄₅ → Cys)] | | • | | | | |
| ApoE ₋₁ [ApoE-2 (Lys ₁₄₆ → Gln)] | | • | | | | |
| ApoE ₋₂ (ApoE _{Bethesda}) | • | | | | | |
| ApoE ₋₂ [ApoE-1 (Gly ₁₂₇ → Asp, Arg ₁₅₈ → Cys)] | • | | | | | |
| ApoE ₊₂ (ApoE-5) | | | | | • | |
| ApoE Absence | | | | | | |

standardized designation of apoE isoforms based on unit charge with an accompanying designation in parentheses as the sequence and (or) post-translational modifications of the isoform become known.

ApoH

The glycoprotein ApoH (*M_r* 46 000) is also designated β_2 -glycoprotein-I (91, 92). Reportedly, it modulates the activity of lipoprotein lipase in vitro (92). Identification of sialylated apoH on the electrophoretogram of plasma is difficult because of the streaking of the albumin bands (Figure 1). Treatment of apoH with neuraminidase shifts the electrophoretic position of apoH to a pI = 9.5 (Figure 14). Desialylated apoH is not visible on the electrophoretogram in Figure 2 because of the narrow pH range.

Discussion

The analytical system we describe is sensitive and reproducible for evaluating apolipoproteins in normal subjects and patients with dyslipoproteinemias. Variants, as well as deficiencies of individual apolipoproteins, can now be readily identified in human plasma. This procedure can also be used to study the biosynthesis of apolipoproteins by cells, in addition to intracellular and extracellular processing and modification of apolipoproteins. Of particular interest is an evaluation of the coordinated control of the biosynthesis of several different apolipoproteins and the factors that modulate lipoprotein biosynthesis.

By using a narrow pH range (pH 4–7) and a sensitive silver stain, we have enhanced the resolution of apolipoproteins. Complications induced by protein-protein interactions via disulfide bonds are eliminated by reducing the sample before the analysis. (Mixed disulfides are particular-

ly important in apoE-apoA-II dimers and apoE-apoE dimers.) Reduction of the sample also significantly increases the quantity of monomer apoA-II and apoE.

In the present study, we identified two new proteins on the plasma two-dimensional gel electrophoretograms: apoD and apoH. Both are glycoproteins, with many polymorphic isoforms present. Most of the isoforms are eliminated by treatment with neuraminidase.

Use of computerized gel-image systems for apolipoprotein quantification will provide additional information on plasma apolipoprotein concentrations and isoform distribution in normal subjects and patients with dyslipoproteinemias. This quantification technique is currently under investigation.

We extend our gratitude to Rosemary Ronan for the isolation of the apolipoproteins, to the NIH graphic arts department and the photography department (particularly R. V. Dreyfuss) for their excellent technical assistance, and to Imogene Surrey for typing this manuscript.


References

1. Brewer HB Jr, Zech LA, Gregg RE, et al. Type III hyperlipoproteinemia: Diagnosis, molecular defects, pathology, and treatment. *Ann Intern Med* 98, 623–640 (1983).
2. Avogaro P, Cazzolato G, Bon GB, Quinci GB. Are apolipoproteins better discriminators than lipids for atherosclerosis? *Lancet* i, 901–903 (1979).
3. Sniderman A, Shapiro S, Marpole D, et al. Association of coronary atherosclerosis with hyperapobetalipoproteinemia (increased protein but normal cholesterol levels in human plasma low density (B) lipoproteins). *Proc Natl Acad Sci USA* 77, 604–608 (1980).
4. Wintrobe MM, Lee GR, Boggs DR, et al., Eds. The abnormal hemoglobins: General principles. In *Clinical Hematology*, 8th ed., Chap. 33, Lea and Febiger, Philadelphia, PA, 1981, pp 804–816.
5. Harris H. *The Principles of Human Biochemical Genetics*, 3rd ed., Elsevier North Holland Press, New York, NY, 1980, pp 316–405.
6. Zannis VI, Breslow JL, Utermann G, et al. Proposed nomenclature of apoE isoproteins, apoE genotypes, and phenotypes. *J Lipid Res* 23, 911–914 (1982).
7. Zannis VI, Breslow JL, Katz AJ. Isoproteins of human apolipoprotein A-I demonstrated in plasma and intestinal organ culture. *J Biol Chem* 255, 8612–8617 (1980).
8. Utermann G. Human apolipoprotein mutants. In *Atherosclerosis VI*. G Schettler et al., Eds., Springer Verlag, Berlin, 1983, pp 611–615.
9. Anderson NL, Anderson NG. High resolution two-dimensional electrophoresis of human plasma proteins. *Proc Natl Acad Sci USA* 74, 5421–5424 (1977).
10. Anderson NL, Hickman BJ. Analytical techniques for cell

ApoH



ApoH (neuraminidase treated)



pI=3

pI=10

Fig. 14. Two-dimensional electrophoretograms of purified apoH with and without neuraminidase treatment (pH 3–10)

The desialylated form has a pI = 9.5

fractions. XXIV. Isoelectric point standards for two-dimensional electrophoresis. *Anal Biochem* 93, 312-320 (1979).

11. Anderson L, Anderson N. Some perspectives on two-dimensional protein mapping. *Clin Chem* 30, 1898-1905 (1984).

12. Havel RJ, Eder HA, Bragdon JH. The distribution and chemical composition of ultracentrifugally separated lipoproteins in human serum. *J Clin Invest* 34, 1345-53 (1955).

13. Folch J, Lees M, Sloan-Stanley GH. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 226, 495-509 (1957).

14. O'Farrell PH. High resolution two-dimensional electrophoresis of proteins. *Ibid.* 250, 4007-4021 (1975).

15. Zannis VI, Breslow JL. Human very low density lipoprotein apolipoprotein E isoprotein polymorphism is explained by genetic variation and post-translational modification. *Biochemistry* 20, 1033-1041 (1981).

16. Ghiselli G, Gregg RE, Brewer HB Jr. Apolipoprotein E^{Bethesda}. Apolipoprotein E isolated from very low density lipoproteins. *Biochim Biophys Acta* 794, 333-339 (1984).

17. Merrill CR, Goldman D, Sedman SA, Ebert MH. Ultrasensitive stain for proteins in polyacrylamide gels show regional variation in cerebrospinal fluid proteins. *Science* 211, 1437-1438 (1981).

18. Morrissey JH. Silver stain for proteins in polyacrylamide gels: A modified procedure with enhanced uniform sensitivity. *Anal Biochem* 117, 307-310 (1981).

19. Towbin H, Staehlin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets—procedure and some applications. *Proc Natl Acad Sci USA* 76, 4350-4354 (1979).

20. Brewer HB Jr, Lux SE, Ronan R, John KM. Amino acid sequence of human apoLp-Glu-II (apoA-II), an apolipoprotein isolated from the high density lipoprotein complex. *Ibid.*, 69, 1304-1308 (1972).

21. McConathy WJ, Alaupovic P. Studies on the isolation and partial characterization of apolipoprotein D and lipoprotein D of human plasma. *Biochemistry* 15, 515-520 (1976).

22. Law SW, Gray G, Brewer HB Jr. cDNA cloning of human apoA-I amino-acid sequence of preproapoA-I. *Biochem Biophys Res Commun* 112, 257-264 (1983).

23. Shoulders CC, Kornblihtf AR, Munro BS, Baralle FE. Gene structure of human apolipoprotein A-I. *Nucl Acid Res* 11, 2827-2837 (1983).

24. Karathanasis SK, Zannis VI, Breslow JL. Isolation and characterization of the human apolipoprotein A-I gene. *Proc Natl Acad Sci USA* 80, 6147-6151 (1983).

25. Law S, Brewer HB Jr. Nucleotide sequence and the encoded amino acids of human apolipoprotein A-I mRNA. *Ibid.*, 81, 66-70 (1984).

26. Brewer HB Jr, Fairwell T, Kay L, et al. Human plasma preapoA-I: Isolation and amino-terminal sequence. *Biochem Biophys Res Commun* 113, 626-632 (1983).

27. Brewer HB Jr, Fairwell T, Larue A, et al. The amino acid sequence of human apoA-I: An apolipoprotein isolated from high density lipoproteins. *Ibid.* 80, 623-630 (1978).

28. Fielding CJ, Shore VG, Fielding PE. A protein cofactor of lecithin:cholesterol acyltransferase. *Ibid.* 46, 1493-1498 (1972).

29. Law SW, Gray G, Brewer HB Jr, et al. Human apolipoprotein A-I and C-III reside in the p11-q13 region of chromosome 11. *Ibid.* 118, 934-942 (1984).

30. Maciejko JJ, Holmes DR, Kottke BA, et al. Apolipoprotein A-I as a marker of angiographically assessed coronary-artery disease. *N Engl J Med* 309, 385-389 (1983).

31. Utermann G, Feussner G, Franceschini G, et al. Genetic variants of group A apolipoproteins. Rapid methods for screening and characterization without ultracentrifugation. *J Biol Chem* 257, 501-507 (1982).

32. Schaefer EJ, Kay LL, Zech LA, Brewer HB Jr. Tangier disease: High density lipoprotein deficiency due to defective metabolism of an abnormal apolipoprotein A-I (apoA-I_{Tangier}). *J Clin Invest* 70, 934-945 (1982).

33. Assmann G. Tangier disease and the possible role of high-density lipoproteins in atherosclerosis. *Atherosclerosis Rev* 6, 1-28 (1979).

34. Menzel HJ, Kladetzky, Assmann G. One-step screening method

for the polymorphism of apolipoproteins A-I, A-II, and A-IV. *J Lipid Res* 23, 915-922 (1982).

35. Weisgraber KH, Bersot TP, Mahley RW, et al. A-I_{Milano} apoprotein: Isolation and characterization of a cysteine-containing variant of the A-I apoprotein from human high density lipoproteins. *J Clin Invest* 66, 901-907 (1980).

36. Norum RA, Lakier JB, Goldstein S, et al. Familial deficiency of apolipoproteins A-I and C-III and precocious coronary-artery disease. *N Engl J Med* 306, 1513-1519 (1982).

37. Lackner KJ, Law SW, Brewer HB Jr. Human apolipoprotein A-II: Complete nucleic acid sequence of preproapoA-II. *FEBS Lett* 175, 159-164 (1984).

38. Gordon JL, Budelier KA, Sims HF, et al. Biosynthesis of human preproapolipoprotein A-II. *J Biol Chem* 258, 14054-14059 (1983).

39. Lackner KJ, Law SW, Brewer HB Jr, et al. The human apolipoprotein A-II gene is located on chromosome 1. *Biochem Biophys Res Commun* 122, 877-883 (1984).

40. Jahn CE, Osborne JC Jr, Schaefer EJ, Brewer HB Jr. Activation of the enzymatic activity of hepatic lipase by apolipoprotein A-II. Characterization of a major component of high density lipoprotein as the activation plasma component in vitro. *Eur J Biochem* 131, 25-29 (1983).

41. Albers JJ, Cheung MC, Hazzard WR. High-density lipoproteins in myocardial infarction survivors. *Metabolism* 27, 479-485 (1978).

42. Schmitz G, Ilsemann K, Melnik B, Assmann G. Isoproteins of human apolipoprotein A-II: Isolation and characterization. *J Lipid Res* 24, 1021-1029 (1983).

43. Lackner KJ, Edge SB, Gregg RE, et al. Isoforms of apoA-II in human plasma and thoracic duct lymph: Identification of proapoA-II and sialic acid containing isoforms. Submitted.

44. Fidge N, Nestel P. Studies on the in vivo metabolism of A-IV apoprotein in human subjects. *Arteriosclerosis* 1, 377 (1981). Abstract.

45. Gordon JI, Smith DP, Alpers DH, Strauss AW. Cloning of a complementary deoxyribonucleic-acid encoding a portion of rat intestinal preapolipoprotein A-IV messenger ribonucleic-acid. *Biochemistry* 21, 5424-5431 (1982).

46. Schamaun O, Olaisen B, Mevag B, et al. Linkage relations of human apoA-I. In Proc. 7th Int. Workshop on Human Gene Mapping. *Cytogenet Cell Genet* 37, 103-126 (1984).

47. Kane JP, Hardman DA, Paulus HE. Heterogeneity of apolipoprotein B: Isolation of a new species from human chylomicrons. *Proc Natl Acad Sci USA* 77, 2365-2369 (1980).

48. Meng MS, Gregg RE, Schaefer EJ, Hoeg JM, Brewer HB Jr. Presence of two forms of apoB in patients with dyslipoproteinemia. *J Lipid Res* 24, 803-809 (1983).

49. Edge N, Hoeg JM, Schneider PD, Brewer HB Jr. Adult human liver synthesis of apolipoprotein B-100 but not apoB-48. *Arteriosclerosis* 3, 499-500 (1983). Abstract.

50. Brown MS, Kovanen PT, Goldstein JL. Regulation of plasma cholesterol by lipoprotein receptors. *Science* 212, 628-635 (1981).

51. Herbert PN, Assmann G, Gotto AM Jr, Fredrickson DS. Familial lipoprotein deficiency: Abetalipoproteinemia, hypobetalipoproteinemia, and Tangier disease. In *The Metabolic Basis of Inherited Disease*, JB Stanbury et al., Eds., McGraw-Hill Book Co., New York, NY, 1983, pp 389-621.

52. Gabelli C, Stark D, Gregg RE, Brewer HB Jr. Separation of apolipoproteins B-100 and B-48 by agarose polyacrylamide gel electrophoresis (manuscript submitted).

53. Kuoff TJ, Tobertson ME, Priestley LM, et al. Characterization of mRNAs encoding the precursor for human apolipoprotein C-I. *Nucl Acid Res* 12, 3909-3915 (1984).

54. Shulman RS, Herbert PN, Wehrly K, Fredrickson DS. The complete amino acid sequence of C-I (apoLp-Ser), an apolipoprotein from human very low density lipoproteins. *J Biol Chem* 250, 182-190 (1975).

55. Jackson RL, Sparrow JT, Baker HN, et al. The primary structure of apolipoprotein-serine. *Ibid.* 249, 5308-5313 (1974).

56. Soutar AK, Garner CW, Baker HN, et al. Effect of human plasma apolipoproteins and phosphatidylcholine acyl donor on activity of lecithin-cholesterol acyltransferase. *Biochemistry* 14, 3057-3064 (1975).

57. LaRosa JC, Levy RI, Herbert PN, et al. A specific apoprotein

- activator for lipoprotein lipase. *Biochem Biophys Res Commun* 41, 57-62 (1970).
58. Havel RJ, Shore VG, Shore B, Bier DM. Role of specific glycopeptides of human serum lipoproteins in the activation of lipoprotein lipase. *Circ Res* 27, 595-600 (1970).
59. Fojo SS, Law SW, Brewer HB Jr. Human apolipoprotein C-II: Complete nucleic acid sequence of preapoC-II. *Proc Natl Acad Sci USA* 81, 6354-6357 (1984).
60. Jackson RL, Baker HN, Gilliam EB, Gotto AM. Primary structure of very low-density apolipoprotein-C-2 of human plasma. *Ibid.* 74, 1942-1945 (1977).
61. Hospattankar AV, Fairwell T, Ronan R, Brewer HB Jr. Amino acid sequence of human plasma apolipoprotein C-II from normal and hyperlipoproteinemic subjects. *J Biol Chem* 259, 318-322 (1984).
62. Jackson CL, Bruns GAP, Breslow JL. Isolation and sequence of human apolipoprotein CII cDNA clone and its use to isolate and map to human chromosome 19 the gene for apolipoprotein CII. *Proc Natl Acad Sci USA* 81, 2945-2949 (1984).
63. Fojo SS, Law SW, Brewer HB Jr, et al. The localization of the gene for apolipoprotein C-II to chromosome 19. *Biochem Biophys Res Commun* 122, 687-693 (1984).
64. Havel RJ, Kotite L, Kane JP. Isoelectric heterogeneity of the cofactor protein for lipoprotein lipase in human blood plasma. *Biochem Med* 21, 121-138 (1979).
65. Breckenridge WC, Little JA, Steiner G, et al. Hypertriglyceridemia associated with deficiency of apolipoprotein C-II. *N Engl J Med* 298, 1265-1273 (1978).
66. Miller NE, Rao SN, Alaupovic P, et al. Familial apolipoprotein C-II deficiency. Plasma-lipoproteins and apolipoproteins in heterozygous and homozygous subjects and the effects of plasma infusion. *Eur J Clin Invest* 11, 69-76 (1981).
67. Crepaldi G, Fellin R, Baggio G, et al. In *Atherosclerosis V*, AM Gotto Jr, LC Smith, B Allen, Eds., Springer-Verlag, New York, NY, 1980, pp 250-254.
68. Sharpe CR, Sidoli A, Shelley CS, et al. Human apolipoproteins AI, AII, CII and CIII. cDNA sequences and mRNA abundance. *Nucl Acid Res* 12, 3917-3932 (1984).
69. Brewer HB Jr, Shulman R, Herbert P, et al. The complete amino acid sequence of alanine apolipoprotein (apoC-III), an apolipoprotein from human plasma very low density lipoproteins. *J Biol Chem* 249, 4975-4984 (1974).
70. Quarfordt SH, Michalopoulos G, Schirmer B. The effect of human C apolipoproteins in the in vitro hepatic metabolism of triglyceride emulsions in the rat. *Ibid.* 257, 1442-1447 (1982).
71. Windler E, Chao YS, Havel RJ. Determinants of hepatic uptake of triglyceride-rich lipoproteins and their remnants in the rat. *Ibid.* 255, 5475-5480 (1984).
72. Schaefer EJ, Law SW, Ghiselli G, et al. Biochemical, clinical, and genetic features in familial apolipoprotein A-I and C-III absence: A new disease associated with high density lipoprotein deficiency and premature coronary artery disease. (manuscript in preparation for publication).
73. Karathanasis SK, Norum RA, Zannis VI, Breslow JL. An inherited polymorphism in the human apolipoprotein A-I gene locus related to the development of atherosclerosis. *Nature (London)* 301, 718-720 (1983).
74. McConathy WJ, Alaupovic P. Isolation and partial characterization of apolipoprotein D: A new protein moiety of the human plasma lipoprotein system. *FEBS Lett* 37, 178-182 (1973).
75. Fielding PE, Fielding CJ. Evidence for a cholesteryl ester transfer complex in human plasma. *Proc Natl Acad Sci USA* 77, 3327-3330 (1980).
76. Zannis VI, McPherson J, Goldbager G, et al. Synthesis, intracellular processing and signal peptide of human apolipoprotein E. *J Biol Chem* 259, 5495-5499 (1984).
77. Mclean JW, Fukazawa C, Taylor JM. Rat apolipoprotein E mRNA. Cloning and sequencing of double stranded cDNA. *J Biol Chem* 258, 8993-9000 (1983).
78. Olaison O, Teisberg P, Gedde-Dahl T Jr. The locus for apolipoprotein E (apoE) is linked to the complement component C3 (C3) locus on chromosome 19 in man. *Hum Genet* 62, 233-236 (1982).
79. Mahley RW, Hui DY, Innerarity TL, Weisgraber KH. Two independent lipoprotein receptors in hepatic membranes of dog, swine, and man. *J Clin Invest* 68, 1197-1206 (1981).
80. Utermann G, Jaeschke M, Menzel J. Familial hyperlipoproteinemia type III: Deficiency of a specific apolipoprotein (apoE-III) in very low density lipoproteins. *FEBS Lett* 56, 352-355 (1975).
81. Gregg RE, Zech LA, Schaefer EJ, Brewer HB Jr. Type III hyperlipoproteinemia: Defective metabolism of an abnormal apolipoprotein E. *Science* 211, 584-586 (1981).
82. Schneider WJ, Kovanen PT, Brown MS, et al. Familial dysbetalipoproteinemia: Abnormal binding of mutant apolipoprotein E to lipoprotein receptors of human fibroblasts and membranes from liver and adrenal of rats, rabbits, and cows. *J Clin Invest* 68, 1075-1085 (1981).
83. Weisgraber KH, Rall SC Jr, Mahley RW. Human E apoprotein heterogeneity. Cysteine-arginine interchanges in the amino acid sequence of the apo-E isoforms. *J Biol Chem* 256, 9077-9083 (1981).
84. Rall SC Jr, Weisgraber KH, Mahley RW. Human apolipoprotein E: The complete amino acid sequence. *Ibid.* 257, 4171-4178 (1982).
85. Rall SC Jr, Weisgraber KH, Innerarity TL, Mahley RW. Structural basis for receptor binding heterogeneity of apolipoprotein E from type III hyperlipoproteinemic subjects. *Proc Natl Acad Sci USA* 79, 4696-4700 (1982).
86. Rall SC Jr, Weisgraber KH, Innerarity TL, et al. Identification of a new structural variant of human apolipoprotein E, E2(Lys146-Gln), in a type III hyperlipoproteinemic subject with the E3/2 phenotype. *J Clin Invest* 72, 1288-1297 (1983).
87. Gregg RE, Ghiselli G, Brewer HB Jr. Apolipoprotein E (Bethesda): A new variant of apolipoprotein E associated with type III hyperlipoproteinemia. *J Clin Endocrinol Metab* 57, 968-974 (1983).
88. Weisgraber KH, Rall SC Jr, Innerarity TL, et al. A novel electrophoretic variant of human apolipoprotein E. Identification and characterization of apolipoprotein E1. *J Clin Invest* 73, 1024-1033 (1984).
89. Yamamura T, Yamamoto A, Hiramori K, Nambu S. A new isoform of apolipoprotein E-apoE-5-associated with hyperlipidemia and atherosclerosis. *Atherosclerosis* 50, 159-172 (1984).
90. Ghiselli G, Schaefer EJ, Gascon P, Brewer HB Jr. Type III hyperlipoproteinemia associated with apolipoprotein E deficiency. *Science* 214, 1239-1241 (1981).
91. Polz E, Kostner GM. Binding of β_2 -glycoprotein-I to Intralipid: Determination of the dissociation constant. *Biochem Biophys Res Commun* 90, 1305-1312 (1979).
92. Nakaya Y, Schaefer EJ, Brewer HB Jr. Activation of human post-heparin lipoprotein lipase by apolipoprotein H (β_2 -glycoprotein-I). *Ibid.* 95, 1168-1172 (1980).