

Selective Removal of Aberrant Extender Units by a Type II Thioesterase for Efficient FR-008/Candidicin Biosynthesis in *Streptomyces* sp. Strain FR-008^{∇†}

Yongjun Zhou,¹ Qingqing Meng,² Delin You,¹ Jialiang Li,¹ Shi Chen,¹ Dazhong Ding,² Xiufen Zhou,¹ Huchen Zhou,^{2*} Linquan Bai,^{1*} and Zixin Deng¹

Laboratory of Microbial Metabolism, School of Life Science & Biotechnology, Shanghai Jiaotong University, Shanghai 200030, China,¹ and School of Pharmacy, Shanghai Jiaotong University, Shanghai 200240, China²

Received 6 May 2008/Accepted 26 September 2008

Gene *fscTE*, encoding a putative type II thioesterase (TEII), was associated with the FR-008/candidicin gene cluster. Deletion of *fscTE* reduced approximately 90% of the FR-008/candidicin production, while the production level was well restored when *fscTE* was added back to the mutant *in trans*. FscTE was unable to compensate for the release of the maturely elongated polyketide as site-directed inactivation of the type I thioesterase (TEI) totally abolished FR-008/candidicin production. Direct biochemical analysis of FscTE in parallel with its homologue TyIO from the tylosin biosynthetic pathway demonstrated their remarkable preferences for acylthioesters (i.e., propionyl-*S*-*N*-acetylcysteamine [SNAC] over methylmalonyl-SNAC and acetyl-SNAC over malonyl-SNAC) and thus concluded that TEII could maintain effective polyketide biosynthesis by selectively removing the nonelongatable residues bound to acyl carrier proteins. Overexpression of FscTE under the strong constitutive *ermE***p* promoter in the wild-type strain did not suppress FR-008/candidicin formation, which confirmed its substrate specificity *in vivo*. Furthermore, successful complementation of the *fscTE* mutant was obtained with *fscTE* and *tyIO*, whereas no complementation was detected with nonribosomal peptide synthetase (NRPS) TEII *tycF* and *srfAD*, reflecting substrate specificities of TEIIs distinctive from those of either polyketide synthases or NRPSs.

Complex polyketides are a large family of bacterial natural products possessing a wide range of biological activities. The carbon frameworks of these compounds are assembled by a common mechanism of decarboxylative condensations of simple malonate derivatives (e.g., malonyl or methylmalonyl) by polyketide synthases (PKSs) in a manner very similar to fatty acid biosynthesis (14). Type I PKSs are complexes of large multimodular enzymes that catalyze biosynthesis of polyketide compounds via repetitive reaction sequences, during which each step is catalyzed by a separate enzymatic domain (7, 15, 33). The biosynthesis logic of type I PKS is also shared by nonribosomal peptide synthetases (NRPSs), which are organized into coordinated modules minimally consisting of adenylation (A), peptidyl carrier protein (PCP), and condensation (C) domains required for an elongation cycle in assembly line arrays (8). The fully extended polyketide or oligopeptide chains bound to terminal enzymatic templates as acyl-acyl carrier protein (ACP) or aminoacyl-PCP thioesters are usually released and cyclized by a type I thioesterase (TEI) domain fused to the carboxyl terminus of the last elongation module (11, 34). Although TEIs have been verified to be sufficient for

chain release and cyclization by many *in vitro* analysis (11, 19, 32, 34), additional thioesterase genes encoding discrete proteins called type II thioesterases (TEIIs) were also found within many type I PKSs and NRPS gene clusters (3–6, 16, 20, 21, 30, 41). Sequence analysis has revealed TEIIs as belonging to the α/β -hydrolase superfamily with a catalytic triad consisting of Ser-Asp-His (21).

TEIIs are generally believed to function as an editing enzyme to restore the biosynthetic machinery by hydrolytically removing the aberrant acyl groups blocking ACP or PCP for further elongation procedures (13, 18, 31). Disruption of the TEII genes greatly reduced the productivity of several antibiotics (3, 6, 30). Coexpression of cognate TEII with PKS in heterologous hosts evidently enhanced polyketide production: e.g., the improved biosyntheses of erythromycin and picromycin aglycones in recombinant strains (27, 37). As the first biochemical evidence of TEII in type I PKS, TyIO from tylosin biosynthesis hydrolyzed the acyl-*N*-acetylcysteamine (acyl-NAC) thioesters simulating the aberrant residues bound to ACP (13). PikAV TEII from picromycin biosynthesis was also tested for its hydrolytic activities with either elongatable or nonelongatable residues attached to ACPs (18). However, PikAV is deficiently correlated with picromycin biosynthesis efficiency as deletion of the TEII gene did not reduce product yield (5).

FR-008/candidicin, a heptaene macrolide antifungal agent, was synthesized by a type I PKS pathway in *Streptomyces* sp. strain FR-008 (Fig. 1) (4, 15, 43). One representative TEII encoded by *fscTE* associated with the FR-008/candidicin (*fsc*) gene cluster shows 48% identity, on the amino acid level, to the TEII (TyIO) from the tylosin biosynthetic pathway (13). In

* Corresponding author. Mailing address for L. Bai: Laboratory of Microbial Metabolism, School of Life Science & Biotechnology, Shanghai Jiaotong University, Shanghai 200030, China. Phone and fax: 86 21 62932418. E-mail: bailq@sjtu.edu.cn. Mailing address for H. Zhou: School of Pharmacy, Shanghai Jiaotong University, Shanghai 200240, China. Phone: 86 21 34206721. Fax: 86 21 34204457. E-mail: hczhou@sjtu.edu.cn.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

[∇] Published ahead of print on 3 October 2008.




FIG. 1. Molecular structure of FR-008/candicidin (A) and the gene cluster associated with FR-008/candicidin biosynthesis (B). A TEII encoded by *fscTE* and a TEI encoded by the C-terminal region of *fscF* are marked by stars.

order to elucidate the substrate specificities of TEII for its editing role in PKS, systematic analyses of FscTE and its homologue TyIO were performed through in vivo gene inactivation followed by complementation and through in vitro catalysis assays with synthesized acyl-thioesters simulating the aberrant or normal extender units in polyketide elongation processes. As an extended insight into the editing role of TEII in PKS, evidence presented here strongly supports distinct substrate specificities of TEII with a capability of selectively removing the nonelongatable residues from the unprocessed PKS proteins, and thus ensuring efficient polyketide biosynthesis.

MATERIALS AND METHODS

Bacterial strains, plasmids, culture conditions, and general techniques. *Streptomyces* sp. strain FR-008 is the wild-type producer of FR-008/candicidins (15). *Escherichia coli* DH10B was used as cloning host. pBluescript II SK(+) (35), pIJ2925 (17), and PMD 18-T vector (TaKaRa) were used for plasmid constructions. pHZ1358 (36) was used for gene replacement in *Streptomyces* sp. strain FR-008, while pIB139 (39)—a pSET152 derivative with the *ermE***p* promoter (2) and a polylinker—was used for mutant complementation. SFM medium (2% agar, 2% mannitol, 2% soybean powder [pH 7.4]) was used for sporulation, fermentation, and conjugation. YEME (0.3% yeast extract, 0.5% peptone, 0.3% malt extract, 1% glucose, 0.3% sucrose [pH 7.2]) liquid medium was used for fermentation (30°C, 2 days). Trypticase soy broth supplemented with 10.3% sucrose and 1% yeast extract was used for growth of mycelia for isolation of total DNA. LB medium was used for *E. coli* propagation. Recombinant DNA techniques were described by Sambrook et al. (29). PCRs were performed using KOD-Plus (Toyobo) or *Taq* DNA polymerase.

Fermentation and analysis of polyene macrolide titer. Polyene samples were extracted with methanol from the spores harvested from agar plates (6 days at 30°C) or the mycelia harvested from liquid fermentation (2 days at 30°C). The antibiotic titer was evaluated through high-performance liquid chromatography (HPLC) analysis at 380 nm. Component FR-008-III was used as the standard for titer comparisons because the abundance ratio of the three main FR-008/candicidin components is almost invariant in the extracts of the wild-type and recombinant strains.

Liquid chromatography-mass spectrometry (LC-MS) analysis was performed using the Agilent 1100 series LC/MSD Trap system. An Agilent Eclipse XDB-C₁₈ column with dimensions of 4.6 by 250 mm was used, and the mobile phase was 45% CH₃CN in 5.5 mM ammonium acetate (pH 4.5) at a flow rate of 0.6 ml/min. The ion trap mass spectrometer was operated with the electrospray ionization source in negative or positive mode. Drying gas flow was 10 liters/min, and nebulizer pressure was 50 lb/in². Drying gas temperature was 350°C. The fragmentation amplitude varied between 1.0 and 1.8 V.

Plasmid construction. Plasmid construction is described in the supplemental material.

Cloning, expression, and purification of FscTE and TyIO. A 929-bp NdeI-EcoRI DNA fragment containing *fscTE* was excised from pJTU598 and cloned into the expression vector pET28a digested with NdeI and EcoRI to generate pJTU2224. pJTU2224 was transformed into *E. coli* BL21(DE3)/pLysE for induced expression of FscTE. pMLH27 (13) was also transformed into *E. coli* BL21(DE3)/pLysE for expression of the previously described TyIO. A 1-liter culture of each recombinant strain was grown using LB medium containing kanamycin (25 mg/ml) and chloramphenicol (12.5 mg/ml). The expression of the N-terminal His₆-tagged recombinant proteins was induced at an optical density at 600 nm of 0.8 with 0.2 mM isopropyl-β-D-thiogalactopyranoside (IPTG), and incubation was continued for 20 h at 18°C.

The cells from each 1-liter culture were harvested by centrifugation at 5,000 × g for 10 min, resuspended in 25 ml of buffer A (50 mM Tris, 300 mM NaCl [pH 7.4]), and lysed by ultrasonication with a Sanyo Soniprep 150. Cell debris was removed by ultracentrifugation (twice for 30 min each at 15,000 × g), and the supernatant was loaded onto a 5-ml Hitrap chelating HP column (Amersham Biosciences). Protein purification procedures were performed at 4°C with an ÄKTA fast protein liquid chromatograph. The flow rate was 0.5 ml/min, and the absorbance was monitored at 280 nm. The column was washed with buffer A followed by a linear gradient of buffer B (50 mM Tris, 300 mM NaCl, 500 mM imidazole [pH 7.4]): from 0 to 100% buffer B over 20 min and 100% buffer B for 5 min. For the activity assay, the protein was concentrated and exchanged into 100 mM phosphate buffer (pH 7.4) with the Amicon Ultra-15 (Millipore) and mixed with 20% glycerol for storage at -80°C. Protein concentrations were determined by Bradford assay (Bio-Rad). Typically, 1 liter of culture yielded 8.8 mg of FscTE and 6.2 mg of TyIO.

Hydrolytic activity assay of TEII. Hydrolysis of acyl-S-N-acetylcysteamine (acyl-SNAC) thioesters releases thiol, which reacts with 5,5'-dithio-2-nitrobenzoic acid (DTNB) to form the detectable chromophore 5-thio-2-nitrobenzoate (λ_{max} , 412 nm; ϵ , 13,600/M/cm) by spectrophotometer (PerkinElmer Lambda

650). To establish the time course of the TEII-catalyzed hydrolysis at 30°C, each assay mixture contained (in a total of 700 μ l) 100 mM sodium phosphate (pH 7.4), 1.24 μ M FscTE or 1.98 μ M TylO, 7 μ l of 20 mM DTNB in 100 mM sodium phosphate (pH 7.4), and various amounts of substrates dissolved in dimethyl sulfoxide. Hydrolysis rates were measured over the concentration range of 10 to 70 mM for NAC thioesters. *p*-Aminobenzoyl-SNAC was assayed only at 5 mM. Reactions were carried out in duplicate, and the hydrolysis rates were calculated from the initial linear portion of the curves. In all cases, the rates were calibrated with the background hydrolysis without added enzyme.

Compound synthesis. Reagents for assays and for chemical synthesis were purchased from Sigma-Aldrich Chemical Co., Ltd. The details of the procedures for synthesis of NAC thioester derivatives are described in the supplemental material. The prepared compounds were further separated on a silica gel column impregnated with copper sulfate (10) or by preparative reverse-phase HPLC (Shimadzu) on a Shimadzu C₁₈ 20- by 250-mm column at a flow rate of 5 ml/min using the mobile phase of 35% CH₃CN-H₂O. Nuclear magnetic resonance spectra for protons (¹H NMR) were recorded on a Varian Mercury 400-MHz spectrometer.

RESULTS

Inactivation of *fscTE* encoding TEII drastically reduced FR-008/candicidin production. To assess the role of *fscTE* in FR-008/candicidin biosynthesis, a 675-bp internal DNA fragment coding for 225 amino acids (aa) of FscTE (285 aa) was in-frame deleted, which contains the entire catalytic triad assumed to be essential for thioester hydrolysis (Fig. 2A). The conjugation plasmid pJTU585, carrying a fragment with the 675-bp deletion, was introduced into *Streptomyces* sp. strain FR-008 by conjugation. After initial selection for thiostrepton resistance (Thio^r) exconjugants followed by two rounds of growth in the absence of thiostrepton, 1 Thio^s derivative out of 20 was selected as a desired mutant, named ZYJ-4, by PCR screening with the primers New-*fscTE* Test-S (5'-TCGGGCG TCCTGCTGCTCCTGCTG-3') and New-*fscTE* Test-A (5'-T CGTCGTTGCGGATGACCTCGGG-3') (Fig. 2B). ZYJ-4 was finally confirmed by sequencing the PCR product amplified from it. About 10% of the original FR-008/candicidin production of the wild-type strain could be detected in ZYJ-4 by LC-MS analysis (Fig. 2C). The necessity of FscTE for the effective FR-008/candicidin biosynthesis was further supported by the evidence that the antibiotic titer was well restored (75%) after complementation of the *fscTE* mutant in *trans* (Fig. 3A). This was achieved by placing the PCR-amplified *fscTE* gene under the direct control of the strong constitutive *ermE***p* promoter on an integrative vector of pJTU598, which was introduced into ZYJ-4 by conjugation. Furthermore, pJTU2216, as a control that differed from pJTU598 with a mutated *fscTE* (S129A), was also introduced into ZYJ-4 and provided no contribution to the restoration of FR-008/candicidin titer (Fig. 3A).

Comparison of FR-008/candicidin productivity with various levels of *fscTE* expression. In order to test whether overexpression of FscTE suppresses FR-008/candicidin biosynthesis by overactive cleavage of the correct residues (e.g., malonyl group) from ACPs (18), we decided to assess FR-008/candicidin biosynthesis efficiency by changing the expression level of *fscTE*. *fscTE* was respectively placed and controlled under the promoters of PKS gene *fscD*, a post-PKS modification gene *fscMI* probably cotranscribed with *fscTE*, or a strong constitutive *ermE***p* promoter. These constructs were individually introduced into the *fscTE* mutant and the wild type as well. In the *fscTE* mutant, *ermE***p*-controlled expression of *fscTE* restored




FIG. 2. Inactivation of TEII gene *fscTE* in *Streptomyces* sp. strain FR-008. (A) Schematic in-frame deletion of 225 aa in FscTE to generate *fscTE* mutant ZYJ-4. The expected PCR product from the wild-type (WT) strain is 1,921 bp, and that from ZYJ-4 is 1,246 bp with the primers New-*fscTE* Test-S and -A. (B) PCR analysis of wild-type and mutant genomic DNAs. PCR products were run on an agarose gel. (C) HPLC analysis of FR-008/candicidin production in wild-type and ZYJ-4. FR-008-V, -III, and -VI are represented by V, III, and VI, respectively. mAU, milliabsorbance units.

FR-008/candicidin titer to 75% of the wild-type titer, much higher than that controlled by *fscMI* or *fscDp* (47% and 49%, respectively). Noticeably, an extra copy of *fscTE* controlled by *ermE***p* did not suppress the FR-008/candicidin titer (72%) compared with that in the wild type carrying the vector pIB139 alone (63%). However, for unknown reasons, the integration of pIB139 into the chromosome of *Streptomyces* sp. strain FR-008 caused 37% reduced FR-008/candicidin production (Fig. 3B).

Restoration of FR-008/candicidin production in the *fscTE* mutant by introduction of different TEIIs. To investigate whether FscTE is catalytically comparable to its homologs from other PKSs and NRPSs, *tylO* (13) was selected to repre-




FIG. 3. (A) FR-008 productions in *fscTE* mutant (ZYJ-4) complemented with different TEIIs. *fscTE* (S129A) represents site-directed mutated TEII; *tylO* is a PKS TEII from the tylosin biosynthetic gene cluster; *tycF* and *srfAD* represent the NRPS TEIIs, respectively, from surfactin and tyrocidine biosynthetic gene clusters. All of the TEII genes were controlled by the *ermE***p* promoter on the integrative vector pIB139. WT, wild type. (B) Comparison of the FR-008/candicidin productivity with various levels of *fscTE* expression. *fscD*p is a promoter controlling the PKS gene *fscD*. *fscMI*p is a promoter of the post-PKS modification gene *fscMI*, which is probably cotranscribed with *fscTE* in one operon.

sent the TEII from PKSs and *tycF* and *srfAD* (40) were selected as the representative TEIIs from NRPSs. The TEII genes were each placed under the control of the constitutive promoter *ermE***p* and then introduced into ZYJ-4. LC-MS analysis of the recombinant strains indicated that *tylO* partially restored the FR-008/candicidin titer to 30%, much lower than the complementation effect of *fscTE*. Moreover, *tycF* and *srfAD* from NRPS biosynthetic gene clusters did not make any contribution to the titer restoration (Fig. 3A).

Inactivation of TEI totally destroyed FR-008/candicidin biosynthesis. Although the release of maturely assembled polyketide is generally believed to be sufficiently carried out by TEI (11, 19, 32, 34), whether TEIIs contribute to this process is not well investigated by in vivo analysis. The active site of the TEI domain in FscF was site-specifically mutated to assess whether FR-008/candicidin was still produced without the catalysis of TEI and, therefore, to explore whether other cognate activity, e.g., TEII, could substitute TEI for the role of terminal release of FR-008/candicidin aglycone. One representative active site triad, comprised of Asp-1845, His-1941, and Ser-1887, was found in the TEI domain at the C terminus of FscF. The Ser residue in the catalytic triad was proved to be essential for the covalent attachment of the maturely assembled chain by structural elucidation and site-directed mutagenesis analysis of TEI (23, 38). For inactivation of TEI in FR-008 strain, the

putative active site Ser-1887 and its neighboring residue, Ser-1888, were both changed into Ala residues and one SacII restriction site was simultaneously introduced to facilitate mutant screening (Fig. 4A). Plasmid pJTU2222, carrying the mutations S1887A and S1888A in *fscF*, was introduced into *Streptomyces* sp. strain FR-008 through conjugation. After initial selection for Thio^r exconjugants followed by two rounds of growth in the absence of thioestrepton, The Thio^s derivatives were screened by a PCR with the primers TE-I-L-S and TE-I-R-A. The desired mutant, named ZYJ-8, was selected by SacII digestion of the PCR products amplified from genomic DNA (Fig. 4B). ZYJ-8 was further confirmed through sequencing of the PCR product. The fermentation extract of ZYJ-8 was analyzed by LC-MS, and no FR-008/candicidin component was detected (Fig. 4C). The result was confirmed by analysis of three parallel mutants. Interestingly, trace amounts of polyene compounds with typical heptaene UV absorption spectra were detected in ZYJ-8, and the one with highest yield and the retention time at 21 min among them was identified with MS spectra of $[M-H]^- = 754.2$.

Overexpression of FscTE and TyIO. FscTE and TyIO were, respectively, expressed from plasmids pJTU2224 and pMLH27 (13) in *E. coli* as N-terminal His₆-tagged recombinant proteins and purified to homogeneity by Ni²⁺ affinity column chromatography. The purified FscTE and TyIO proteins migrated as a single band on sodium dodecyl sulfate-polyacrylamide gel electrophoresis with molecular masses of 35 kDa and 30 kDa (see Fig. S1 in the supplemental material), consistent with their calculated molecular masses of 33,855 Da and 29,848 Da, respectively.

Substrate specificity of FscTE and TyIO. The NAC thioesters were usually used as the substrates in thioesterase hydrolysis assays (11–13, 22, 40). In this study, acetyl-SNAC and propionyl-SNAC were prepared to mimic the aberrant acyl-ACP substrates, whereas malonyl-SNAC and methylmalonyl-SNAC were used to represent the correct substrates in the PKS condensing reactions of FR-008/candicidin and tylosin biosynthesis. On the other hand, *p*-aminobenzoyl-SNAC was also prepared to imitate the *p*-aminobenzoyl-ACP substrate in initiation of FR-008/candicidin biosynthesis (see Table S1 in the supplemental material).

To investigate the substrate recognition of TEII, FscTE and TyIO were incubated with the NAC thioester substrates. Both of FscTE and TyIO showed significant efficiency for hydrolysis of acetyl-SNAC and propionyl-SNAC thioesters (Table 1), whereas very limited hydrolysis was detected with malonyl-SNAC and methylmalonyl-SNAC thioesters (Fig. 5), which excluded the practical possibility of kinetic measurement. Series of time course experiments with substrate concentration ranges of 10 to 70 mM were carried out with consistent substrate preference as presented in Fig. 5. Moreover, *p*-aminobenzoyl-SNAC was not identified as the proper substrate for FscTE and TyIO, which suggests TEII may not interfere the initiation of FR-008/candicidin biosynthesis (data not shown).

DISCUSSION

Some nonelongatable acyl groups, e.g., acetyl or propionyl, were inevitably generated during polyketide biosynthesis. In the absence of a suitable intermediate acyl thioester bound to




FIG. 4. Site-directed mutagenesis of the TEI domain in FscF. (A) Schematic mutagenesis of the TEI domain in FscF to generate mutant ZYJ-8. The putatively essential serines in the catalytic triad were changed to alanines (S1887A and S1888A). One SacII restriction site was introduced to facilitate mutant screening. WT, wild type. (B) SacII digestion of PCR products from the wild-type and mutant strains with the primers TE-I-L-S and TE-I-R-A. (C) HPLC analysis of FR-008/candicidin production in the wild type and ZYJ-8. V, III, and VI disappeared in ZYJ-8. mAU, milliabsorbance units.

the active site thiol, the ketosynthase domain can erroneously decarboxylate the chain extender units such as malonyl or methylmalonyl, resulting in acyl residues blocking ACP from further polyketide elongation procedure (13, 18). Alternatively,

phosphopantetheinyl transferases (PPTases) usually utilize coenzyme A-SH as a 4'-phosphopantetheine donor for activation of apo-ACP/or PCP to the active holo form. However, PPTases occasionally can transfer acyl-phosphopantetheinyl from acyl coenzyme A to ACP/or PCP, generating the unprocessed acyl-ACP/or PCP (28, 31). Reasonably, removal of the nonelongatable acyl groups from ACP through hydrolysis by a thioesterase (e.g., TEII) is necessary to restore PKS function and maximize polyketide production.

Deletion of TEII-encoding genes generally led to a drastic reduction in product yields (3, 6, 9, 30). Consistently, in-frame deletion of the TEII-encoding gene *fscTE* reduced about 90% of FR-008/candicidin production, and then the titer was well restored after complementation of the mutant in *trans*. Addi-

TABLE 1. Kinetic parameters of TEII-catalyzed NAC thioester hydrolysis

Substrate	TEII	k_{cat} (min^{-1})	K_m (mM)	k_{cat}/K_m ($\text{M}^{-1} \text{s}^{-1}$)
Acetyl-SNAC	FscTE	71.6	33.4	35.7
	TyIO	14.8	42.8	5.8
Propionyl-SNAC	FscTE	109.2	32.0	56.9
	TyIO	23.2	35.6	10.9




FIG. 5. Hydrolytic activities of FscTE and TyIO. (A) FscTE with propionyl-SNAC and methylmalonyl-SNAC. (B) FscTE with acetyl-SNAC and malonyl-SNAC. (C) TyIO with propionyl-SNAC and methylmalonyl-SNAC. (D) TyIO with acetyl-SNAC and malonyl-SNAC. The substrate concentration was 20 mM in each assay.

tionally, complementation with site-specific mutated FscTE (S129A) was unable to restore the production, which further confirmed the catalytic activity of TEII is essential for the effective biosynthesis of FR-008/candicidin. Moreover, the fact that there were no FR-008/candicidin components detected in the TEI mutant implied that FscTE cannot substitute for the cognate TEI of the last module for polyketide termination and cyclization.

By selectively removing the nonelongatable residues blocking the ACP from further polyketide elongation, TEII likely catalyzes the regeneration of ACP domains and thus improves polyketide biosynthesis (Fig. 6). The two PKS TEIIs FscTE and TyIO were characterized, in this work, with remarkable hydrolytic activities for the nonelongatable acetyl and propionyl thioesters but very low activities for the correct malonyl and methylmalonyl thioesters required for FR-008/candicidin and tylosin biosyntheses. Correspondingly, another PKS TEII, PikAV, was also demonstrated with a distinct hydrolysis pref-




FIG. 6. FscTE specifically removes the nonelongatable residues (acetyl or propionyl) bound to ACP in PKS elongation steps.

erence for propionyl-ACP (k_{cat}/K_m , $15.8 \pm 1.8 \text{ M}^{-1} \text{ s}^{-1}$) over methylmalonyl-ACP (k_{cat}/K_m , $3.3 \pm 1.1 \text{ M}^{-1} \text{ s}^{-1}$) (18).

Interestingly, PikAV seems to have adverse activity to polyketide biosynthesis as it can cleave the malonyl thioester group from ACP with comparable k_{cat}/K_m value to acetyl thioester group (18). Thus, the adverse activity of TEII was assumed to be responsible for the reduced product yield after overexpression of PikAV TEII. In contrast to the adverse effect of PikAV overexpression, enhanced expression of FscTE did not suppress FR-008/candicidin formation, supporting the notion that TEIIs are only specifically responsible for the removal of aberrant residues from ACP rather than disturbing the normal process. Additionally, the adverse effect of TEII had not been found in NRPS systems so far, where TEIIs were also reported to improve polypeptide biosynthesis by hydrolytically removing the unprocessed acetyl or aminoacetyl residues from misprimed NRPS proteins (31, 40). In the in vitro simulation of NRPS assembly reaction, SrfAD TEII did not suppress the tripeptide formation, and also, no disturbing effect was observed while the YbtT TEII was incubated, at 10-fold molar excess, with the in vitro reaction of a reconstituted yersiniabactin NRPS/PKS assembly line (24, 31). As a futile reaction, TEII-catalyzed hydrolysis of the elongatable substrates should not be the physiological role of TEIIs.

We also compared the restored product yields in the *fscTE* mutant by combining *fscTE* with either the strong constitutive promoter (*ermE*^{*}*p*) or the *fsc* promoters (*fscDp* and *fscMlp*). The constitutive *ermE*^{*}*p* promoter was found to be more efficient than the native promoters *fscDp* and *fscMlp* from FR-008/candicidin gene cluster (Fig. 4B), suggesting that

the constitutive expression of *pscTE* can well maintain the efficiency of FR-008/candicidin biosynthesis.

In accordance with previous studies (13, 18, 31, 40), the two TEIs characterized were also demonstrated with high K_m values (around 33 mM) to acyl thioesters, which implies low affinity of TEIs for acyl-ACP. However, compared to the carboxylated acyl-ACP units processed readily by the PKS, the aberrant residues bound to unprocessed PKS protein with an increased half-life may serve as better substrates for TEI in vivo (13, 18).

The nonequivalent catalysis of TEIs from PKSs and NRPSs has been distinctively presented in the text. TyIO (13) restored 30% productivity of FR-008/candicidin in the *pscTE* mutant, whereas two NRPS TEIs, SrfAD and TycF (40), were null for restoring production. NbmB, the PKS TEI, was also reported to be successful in complementing the *tyIO* mutant for tylosin production (3). In the biochemical assay, SrfAD and BacT, the NRPS TEIs, were exhibited with hydrolytic activity to acetyl-PCP but with no activity to acetyl-ACP (31). In addition, PikAV, the PKS TEI, has comparable activities toward the methylmalonyl thioester bound to the ACPs derived from either PikAIII or DEBS1 (18). The nonequivalent catalysis of TEIs from PKSs and NRPSs might attribute to their recognition to ACP or PCP.

TEI seems to be generally required for maximizing the efficiencies of polyketide or oligopeptide biosynthesis, since the intrinsic errors occurred in PKSs or NRPSs (13, 18, 31, 40) and the TEI genes associated with many of their gene clusters (3–6, 16, 20, 21, 30). The absence of TEI genes in some gene clusters (36, 42) may be compensated for by their homologs associated with other clusters elsewhere in chromosome, as many streptomycetes usually contain dozens of distinct biosynthetic gene clusters (1, 25, 26, 36).

ACKNOWLEDGMENTS

We thank Christopher T. Walsh for providing plasmids pET30a(TycF) and pET30a(srfA-D) and Peter F. Leadlay for providing plasmids pMLH27 and pIB139. We thank David E. Cane for the gift of SNAC thioester samples. We also thank Russell J. Cox for providing the procedure of preparing malonyl-SNAC and Weiguo He for suggestions on methylmalonyl-SNAC synthesis.

This work received financial support from the National Science Foundation of China, the Ministry of Science and Technology (973 and 863 Programs), the Shanghai Municipal Council of Science and Technology, and Shanghai Leading Academic Discipline Project B203.

REFERENCES

- Bentley, S. D., K. F. Chater, A. M. Cerdeno-Tarraga, G. L. Challis, N. R. Thomson, K. D. James, D. E. Harris, M. A. Quail, H. Kieser, D. Harper, A. Bateman, S. Brown, G. Chandra, C. W. Chen, M. Collins, A. Cronin, A. Fraser, A. Goble, J. Hidalgo, T. Hornsby, S. Howarth, C. H. Huang, T. Kieser, L. Larke, L. Murphy, K. Oliver, S. O'Neill, E. Rabinowitz, M. A. Rajandream, K. Rutherford, S. Rutter, K. Seeger, D. Saunders, S. Sharp, R. Squares, S. Squares, K. Taylor, T. Warren, A. Wietzorrek, J. Woodward, B. G. Barrell, J. Parkhill, and D. A. Hopwood. 2002. Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). *Nature* **417**:141–147.
- Bibb, M. J., G. R. Janssen, and J. M. Ward. 1985. Cloning and analysis of the promoter region of the erythromycin resistance gene (*ermE*) of *Streptomyces erythraeus*. *Gene* **38**:215–226.
- Butler, A. R., N. Bate, and E. Cundliffe. 1999. Impact of thioesterase activity on tylosin biosynthesis in *Streptomyces fradiae*. *Chem. Biol.* **6**:287–292.
- Chen, S., X. Huang, X. Zhou, L. Bai, J. He, K. J. Jeong, S. Y. Lee, and Z. Deng. 2003. Organizational and mutational analysis of a complete FR-008/candicidin gene cluster encoding a structurally related polyene complex. *Chem. Biol.* **10**:1065–1076.
- Chen, S., J. B. Roberts, Y. Xue, D. H. Sherman, and K. A. Reynolds. 2001. The *Streptomyces venezuelae* pikAV gene contains a transcription unit essential for expression of enzymes involved in glycosylation of narbonolide and 10-deoxymethanolide. *Gene* **263**:255–264.
- Doi-Katayama, Y., Y. J. Yoon, C. Y. Choi, T. W. Yu, H. G. Floss, and C. R. Hutchinson. 2000. Thioesterases and the premature termination of polyketide chain elongation in rifamycin B biosynthesis by *Amycolatopsis mediterranei* S699. *J. Antibiot. (Tokyo)* **53**:484–495.
- Donadio, S., M. J. Staver, J. B. McAlpine, S. J. Swanson, and L. Katz. 1991. Modular organization of genes required for complex polyketide biosynthesis. *Science* **252**:675–679.
- Finking, R., and M. A. Marahiel. 2004. Biosynthesis of nonribosomal peptides. *Annu. Rev. Microbiol.* **58**:453–488.
- Geoffroy, V. A., J. D. Fetherston, and R. D. Perry. 2000. *Yersinia pestis* YbtU and YbtT are involved in synthesis of the siderophore yersiniabactin but have different effects on regulation. *Infect. Immun.* **68**:4452–4461.
- Gilbert, I. H., M. Ginty, J. A. O'Neill, T. A. Simpson, J. Staunton, and C. L. Willis. 1995. Synthesis of β -keto and α,β -unsaturated N-acetylcysteine thioesters. *Bioorg. Med. Chem.* **5**:1587–1590.
- Gokhale, R. S., D. Hunziker, D. E. Cane, and C. Khosla. 1999. Mechanism and specificity of the terminal thioesterase domain from the erythromycin polyketide synthase. *Chem. Biol.* **6**:117–125.
- Harvey, B. M., H. Hong, M. A. Jones, Z. A. Hughes-Thomas, R. M. Goss, M. L. Heathcote, V. M. Bolanos-Garcia, W. Kroutil, J. Staunton, P. F. Leadlay, and J. B. Spencer. 2006. Evidence that a novel thioesterase is responsible for polyketide chain release during biosynthesis of the polyether ionophore monensin. *ChemBiochem* **7**:1435–1442.
- Heathcote, M. L., J. Staunton, and P. F. Leadlay. 2001. Role of type II thioesterases: evidence for removal of short acyl chains produced by aberrant decarboxylation of chain extender units. *Chem. Biol.* **8**:207–220.
- Hopwood, D. A. 1997. Genetic contributions to understanding polyketide synthases. *Chem. Rev.* **97**:2465–2498.
- Hu, Z., K. Bao, X. Zhou, Q. Zhou, D. A. Hopwood, T. Kieser, and Z. Deng. 1994. Repeated polyketide synthase modules involved in the biosynthesis of a heptaene macrolide by *Streptomyces* sp. FR-008. *Mol. Microbiol.* **14**:163–172.
- Hu, Z., B. A. Pfeifer, E. Chao, S. Murli, J. Kealey, J. R. Carney, G. Ashley, C. Khosla, and C. R. Hutchinson. 2003. A specific role of the *Saccharopolyspora erythraea* thioesterase II gene in the function of modular polyketide synthases. *Microbiology* **149**:2213–2225.
- Janssen, G. R., and M. J. Bibb. 1993. Derivatives of pUC18 that have BglII sites flanking a modified multiple cloning site and that retain the ability to identify recombinant clones by visual screening of *Escherichia coli* colonies. *Gene* **124**:133–134.
- Kim, B. S., T. A. Cropp, B. J. Beck, D. H. Sherman, and K. A. Reynolds. 2002. Biochemical evidence for an editing role of thioesterase II in the biosynthesis of the polyketide pikromycin. *J. Biol. Chem.* **277**:48028–48034.
- Kohli, R. M., J. W. Trauger, D. Schwarzer, M. A. Marahiel, and C. T. Walsh. 2001. Generality of peptide cyclization catalyzed by isolated thioesterase domains of nonribosomal peptide synthetases. *Biochemistry* **40**:7099–7108.
- Kotowska, M., K. Pawlik, A. R. Butler, E. Cundliffe, E. Takano, and K. Kuczek. 2002. Type II thioesterase from *Streptomyces coelicolor* A3(2). *Microbiology* **148**:1777–1783.
- Linne, U., D. Schwarzer, G. N. Schroeder, and M. A. Marahiel. 2004. Mutational analysis of a type II thioesterase associated with nonribosomal peptide synthesis. *Eur. J. Biochem.* **271**:1536–1545.
- Liu, T., D. You, C. Valenzano, Y. Sun, J. Li, Q. Yu, X. Zhou, D. E. Cane, and Z. Deng. 2006. Identification of NanE as the thioesterase for polyether chain release in nanchangmycin biosynthesis. *Chem. Biol.* **13**:945–955.
- Lu, H., S. C. Tsai, C. Khosla, and D. E. Cane. 2002. Expression, site-directed mutagenesis, and steady state kinetic analysis of the terminal thioesterase domain of the methymycin/picromycin polyketide synthase. *Biochemistry* **41**:12590–12597.
- Miller, D. A., L. Luo, N. Hillson, T. A. Keating, and C. T. Walsh. 2002. Yersiniabactin synthetase: a four-protein assembly line producing the nonribosomal peptide/polyketide hybrid siderophore of *Yersinia pestis*. *Chem. Biol.* **9**:333–344.
- Olinyuk, M., M. Samborsky, J. B. Lester, T. Mironenko, N. Scott, S. Dickens, S. F. Haydock, and P. F. Leadlay. 2007. Complete genome sequence of the erythromycin-producing bacterium *Saccharopolyspora erythraea* NRRL23338. *Nat. Biotechnol.* **25**:447–453.
- Omura, S., H. Ikeda, J. Ishikawa, A. Hanamoto, C. Takahashi, M. Shinose, Y. Takahashi, H. Horikawa, H. Nakazawa, T. Osonoe, H. Kikuchi, T. Shiba, Y. Sakaki, and M. Hattori. 2001. Genome sequence of an industrial microorganism *Streptomyces avermitilis*: deducing the ability of producing secondary metabolites. *Proc. Natl. Acad. Sci. USA* **98**:12215–12220.
- Pfeifer, B., Z. Hu, P. Licari, and C. Khosla. 2002. Process and metabolic strategies for improved production of *Escherichia coli*-derived 6-deoxyerythronolide B. *Appl. Environ. Microbiol.* **68**:3287–3292.
- Quadri, L. E., P. H. Weinreb, M. Lei, M. M. Nakano, P. Zuber, and C. T. Walsh. 1998. Characterization of Sfp, a *Bacillus subtilis* phosphopantetheinyl transferase for peptidyl carrier protein domains in peptide synthetases. *Biochemistry* **37**:1585–1595.

29. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
30. Schneider, A., and M. A. Marahiel. 1998. Genetic evidence for a role of thioesterase domains, integrated in or associated with peptide synthetases, in non-ribosomal peptide biosynthesis in *Bacillus subtilis*. *Arch. Microbiol.* **169**:404–410.
31. Schwarzer, D., H. D. Mootz, U. Linne, and M. A. Marahiel. 2002. Regeneration of misprimed nonribosomal peptide synthetases by type II thioesterases. *Proc. Natl. Acad. Sci. USA* **99**:14083–14088.
32. Schwarzer, D., H. D. Mootz, and M. A. Marahiel. 2001. Exploring the impact of different thioesterase domains for the design of hybrid peptide synthetases. *Chem. Biol.* **8**:997–1010.
33. Schwecke, T., J. F. Aparicio, I. Molnar, A. Konig, L. E. Khaw, S. F. Haydock, M. Oliynyk, P. Caffrey, J. Cortes, J. B. Lester, et al. 1995. The biosynthetic gene cluster for the polyketide immunosuppressant rapamycin. *Proc. Natl. Acad. Sci. USA* **92**:7839–7843.
34. Sharma, K. K., and C. N. Boddy. 2007. The thioesterase domain from the pimarinin and erythromycin biosynthetic pathways can catalyze hydrolysis of simple thioester substrates. *Bioorg. Med. Chem. Lett.* **17**:3034–3037.
35. Short, J. M., J. M. Fernandez, J. A. Sorge, and W. D. Huse. 1988. Lambda ZAP: a bacteriophage lambda expression vector with in vivo excision properties. *Nucleic Acids Res.* **16**:7583–7600.
36. Sun, Y., X. Zhou, J. Liu, K. Bao, G. Zhang, G. Tu, T. Kieser, and Z. Deng. 2002. '*Streptomyces nanchangensis*,' a producer of the insecticidal polyether antibiotic nanchangmycin and the antiparasitic macrolide meilingmycin, contains multiple polyketide gene clusters. *Microbiology* **148**:361–371.
37. Tang, L., H. Fu, M. C. Betlach, and R. McDaniel. 1999. Elucidating the mechanism of chain termination switching in the picromycin/methymycin polyketide synthase. *Chem. Biol.* **6**:553–558.
38. Tsai, S. C., L. J. Miercke, J. Krucinski, R. Gokhale, J. C. Chen, P. G. Foster, D. E. Cane, C. Khosla, and R. M. Stroud. 2001. Crystal structure of the macrocycle-forming thioesterase domain of the erythromycin polyketide synthase: versatility from a unique substrate channel. *Proc. Natl. Acad. Sci. USA* **98**:14808–14813.
39. Wilkinson, C. J., Z. A. Hughes-Thomas, C. J. Martin, I. Bohm, T. Mironenko, M. Deacon, M. Wheatcroft, G. Wirtz, J. Staunton, and P. F. Leadlay. 2002. Increasing the efficiency of heterologous promoters in actinomycetes. *J. Mol. Microbiol. Biotechnol.* **4**:417–426.
40. Yeh, E., R. M. Kohli, S. D. Bruner, and C. T. Walsh. 2004. Type II thioesterase restores activity of a NRPS module stalled with an aminoacyl-S-enzyme that cannot be elongated. *ChemBiochem* **5**:1290–1293.
41. Yu, F. M., B. Qiao, F. Zhu, J. C. Wu, and Y. J. Yuan. 2006. Functional analysis of type II thioesterase of *Streptomyces lydicus* AS 4.2501. *Appl. Biochem. Biotechnol.* **135**:145–158.
42. Zhao, C., J. Ju, S. D. Christenson, W. C. Smith, D. Song, X. Zhou, B. Shen, and Z. Deng. 2006. Utilization of the methoxymalonyl-acyl carrier protein biosynthesis locus for cloning the oxazolomycin biosynthetic gene cluster from *Streptomyces albus* JA3453. *J. Bacteriol.* **188**:4142–4147.
43. Zhou, Y., L. J. Zhu, J. Chen, S. Bai, L. Zhou, X. Wu, and H. Z. Deng. 2008. Jun. Incomplete beta-ketone processing as a mechanism for polyene structural variation in the FR-008/candicidin complex. *Chem. Biol.* **15**:629–638.